

Exam 1A

Chem 1142

Spring 2017

Name: _____


MULTIPLE CHOICE. [3 pts ea.] Circle the best response. [45 pts total.]

- Q1. Which ionic compound will likely have the highest melting point?
- a) NaCl
 - b) Na₂S
 - c) MgS
 - d) AlP
 - e) MgCl₂
- Q2. Which substance will have the **greatest** London dispersion forces?
- a) CH₄
 - b) CCl₄
 - c) H₂
 - d) I₂
 - e) Cl₄
- Q3. Which substance will possess dipole-dipole interactions between its molecules?
- a) CO₂
 - b) Br₂
 - c) BF₃
 - d) CF₄
 - e) SO₂
- Q4. Which substance will possess hydrogen-bond interactions between its molecules?
- a) CH₃NH₂
 - b) NF₃
 - c) CH₃OCH₃
 - d) NO₂
 - e) CH₄
- Q5. A cubic unit cell contains tungsten (W) ions at each corner and body, and oxide ions at each face. What is its chemical formula?
- a) W₂O₃
 - b) W₉O₆
 - c) W₃O₂
 - d) WO₂
 - e) W₃O₄

Q6. An example of a network covalent solid is:

- a) brass
- b) ice
- c) quartz
- d) gold
- e) sucrose

Q7. What type of unit cell is shown below:



- a) simple cubic
- b) face-centered cubic
- c) body-centered cubic
- d) tetragonal
- e) orthorhombic

Q8. A solution of NaCl(aq) has a molal concentration of 2.0 m. How many moles of NaCl are present if there are 125-g of $\text{H}_2\text{O(l)}$?

- a) 16-mol
- b) 0.016 mol
- c) 0.25 mol
- d) 63 mol
- e) 2.0 mol


Q9. Which aqueous solution will have the **largest** boiling point? Assume ideal behavior.

- a) 0.100 m glucose
- b) 0.100 m sucrose
- c) 0.300 m lithium phosphate
- d) 0.400 m sodium chloride
- e) 0.500 m ethanol

Q10. A solution of LiCl(aq) has an osmotic pressure of 1.8 atm at a temperature of 35 °C. Calculate the concentration of the solution, assuming ideal behavior.

- a) 0.036 M
- b) 0.071 M
- c) 23 M
- d) 45 M
- e) 0.31 M

- Q11. The boiling point of 1.0 m $\text{FeCl}_3(\text{aq})$ is $101.2\text{ }^\circ\text{C}$. Calculate the van't Hoff factor for FeCl_3 from this data.
Note: $k_b(\text{H}_2\text{O}) = 0.52\text{ }^\circ\text{C}/\text{m}$.
- 4.0
 - 3.8
 - 2.3
 - 1.2
 - 0.60
- Q12. Given the vapor pressure curve for mercury shown below, estimate its normal boiling point.



- 575 K
 - 640 K
 - 400 K
 - 530 K
 - 670 K
- Q13. For most substances, when you are below the critical temperature it is possible to convert the gas phase into either a solid or liquid phase by compression. Above the critical temperature, this is not possible because:
- the substance is a plasma in this region
 - the substance has extremely large IMF in this region
 - the substance is a crystal in this region
 - the substance is a supercritical fluid in this region
 - the substance is volatile in this region
- Q14. Predict which two liquids will likely be miscible:
- CS_2 / C_8H_{18}
 - C_8H_{18} / CH_3OH
 - $\text{CH}_3\text{CH}_2\text{NH}_2$ / C_6H_{14}
 - C_8H_{18} / C_7H_{16}
 - CH_3NH_2 / C_5H_{12}

- Q15. $\text{N}_2(\text{g})$ has a Henry's law constant of $8.2 \times 10^{-4} \text{ M}\cdot\text{atm}^{-1}$ for water at 4°C . If the concentration of N_2 in water is found to be 0.100 M , what must the pressure of $\text{N}_2(\text{g})$ be?
- a) 120 atm
 - b) $8.2 \times 10^{-3} \text{ atm}$
 - c) 0.10 atm
 - d) 0.19 atm
 - e) 10. atm

Short Response.

Show ALL work to receive credit. Be sure to use the conversion-factor (dimensional-analysis) method for all problems involving conversions!

- Q16. [11 pts.] Cesium (Cs) crystallizes in a body-centered cubic unit cell with an edge length of 614.1 pm. Being careful to show all work—including units and significant figures—calculate its density in units of g/cm^3 .

Q17. [11 pts.] i) List the intermolecular forces present between the following molecules:

a) NH_3 _____

b) CH_3F _____

c) CO_2 _____

ii) Explain in detail why MgO has a much greater melting point than SO_2 . Your answer should include complete sentences and diagrams where appropriate.

Q18. [11 pts.] Calculate the freezing point of 12.4 M NaBr(aq), given a solution density of 2.08 g/mL. What assumption are you making? Note, $k_f(\text{H}_2\text{O}) = 1.86 \text{ }^\circ\text{C}/\text{m}$.
Hint: start by converting the molar concentration to a molal concentration!

Q19. [11 pts.] The boiling point of an aqueous solution formed by adding 10.0-g of an unknown non-electrolyte to 150.0-g of water is found to be 100.715 °C. Show how to, and then calculate the molar mass of the unknown substance. Note: $k_b(\text{H}_2\text{O}) = 0.52 \text{ }^\circ\text{C}/\text{m}$.

Q20. [11 pts.] Be sure to show all work!

a) How many moles of NaCl are contained in 325-g of a 1.00 %(w/w) aqueous solution?

b) Water has an unusual pressure-temperature phase diagram with a solid/liquid line that has a negative slope. What does this mean in terms of the melting point as we increase the pressure? You should sketch part of the phase diagram as part of your answer. Your explanation should be in the form of complete sentences.

c) What happens to the following three types of concentrations as the temperature is increased? Be sure to explain your answer.

i) Molal concentration

ii) Molar concentration

iii) Percent by mass, %(w/w)

BONUS Question:

Sketch out the structure of graphite, and explain why it can act as an effective lubricant.



"Rats! I thought lanthanoids and actanoids were gonna be giant robots or something."

Periodic Table of the Elements

IA	IIA										IIIA										IVA	VA	VIA	VIIA	VIIIA		
1																					13	14	15	16	17	18	
1 H 1.01																					5 B 10.81	6 C 12.01	7 N 14.01	8 O 16.00	9 F 19.00	10 Ne 20.18	
3 Li 6.94	4 Be 9.01																					13 Al 26.98	14 Si 28.09	15 P 30.97	16 S 32.07	17 Cl 35.45	18 Ar 39.95
11 Na 22.99	12 Mg 24.31	3	4	5	6	7	8	9	10	11	12	31 Ga 69.72	32 Ge 72.61	33 As 74.92160	34 Se 78.96	35 Br 79.90	36 Kr 83.80										
19 K 39.10	20 Ca 40.08	21 Sc 44.96	22 Ti 47.87	23 V 50.94	24 Cr 52.00	25 Mn 54.94	26 Fe 55.85	27 Co 58.93	28 Ni 58.69	29 Cu 63.55	30 Zn 65.39	49 In 114.82	50 Sn 118.71	51 Sb 121.76	52 Te 127.60	53 I 126.90	54 Xe 131.29										
37 Rb 85.47	38 Sr 87.62	39 Y 88.91	40 Zr 91.22	41 Nb 92.91	42 Mo 95.94	43 Tc [98]	44 Ru 101.07	45 Rh 102.91	46 Pd 106.42	47 Ag 107.87	48 Cd 112.41	81 Tl 204.38	82 Pb 207.20	83 Bi 208.98	84 Po [210]	85 At [210]	86 Rn [222]										
55 Cs 132.91	56 Ba* 137.33	71 Lu 174.97	72 Hf 178.49	73 Ta 180.95	74 W 183.84	75 Re 186.21	76 Os 190.23	77 Ir 192.22	78 Pt 195.08	79 Au 196.97	80 Hg 200.59	113	114	115	116	117	118										
87 Fr [223]	88 Ra** [226]	103 Lr [262]	104 Rf [261]	105 Db [262]	106 Sg [266]	107 Bh [264]	108 Hs [265]	109 Mt [268]	110	111	112																
		* 57 La 138.91	58 Ce 140.12	59 Pr 140.91	60 Nd 144.24	61 Pm [145]	62 Sm 150.36	63 Eu 151.96	64 Gd 157.25	65 Tb 158.93	66 Dy 162.50	67 Ho 164.93	68 Er 167.26	69 Tm 168.93	70 Yb 173.04												
		** 89 Ac [227]	90 Th 232.04	91 Pa 231.04	92 U 238.03	93 Np [237]	94 Pu [244]	95 Am [243]	96 Cm [247]	97 Bk [247]	98 Cf [251]	99 Es [252]	100 Fm [257]	101 Md [258]	102 No [259]												

1 atm = 101,325 Pa = 760 mmHg = 760 torr

$T/K = t/^{\circ}C + 273.15$

$R = 0.08206 \frac{\text{atm} \cdot \text{L}}{\text{mol} \cdot \text{K}}$

$R = 8.3145 \text{ J/mol} \cdot \text{K}$

$\Delta T_b = ik_b m$

$\Delta T_f = ik_f m$

$\Pi = iMRT$

$k_f(\text{H}_2\text{O}) = 1.86 \text{ }^{\circ}\text{C/m}$

$k_b(\text{H}_2\text{O}) = 0.52 \text{ }^{\circ}\text{C/m}$

$c = kP$