

Exam 3A

Chem 1142

Spring 2019



Name: _____

MULTIPLE CHOICE. [2 pts ea.] Record the best response on the scantron sheet. [40 pts total.]


Assume all solutions are aqueous and at a temperature of 25 °C, unless stated otherwise.

- Q1. Bases turn litmus what color?
A) blue
B) green
C) red
D) yellow
- Q2. The Arrhenius definition of an acid is:
A) they donate H^+ ions to other molecules
B) they form H^+ ions in water
C) they accept electron pairs
D) they turn litmus red
- Q3. A good example of an Arrhenius base is:
A) NH_3
B) $LiOH$
C) CH_3CO_2H
D) HCO_3^-
- Q4. The conjugate acid to HPO_4^{2-} is:
A) H_3PO_4
B) $H_2PO_4^-$
C) PO_4^{3-}
D) H_3O^+
- Q5. An example of a strong acid would be:
A) HF
B) HNO_2
C) H_2SO_4
D) H_3PO_4
- Q6. At 37 °C, K_w is equal to 5.5×10^{-14} . This means the pH of pure water at this temperature will be:
A) 7.00
B) 5.50
C) 13.26
D) 6.63


- Q7. Which of the following diagrams represents a snapshot of a very small portion of a beaker containing a weak acid, HA, dissolved in water? Note that the solvent molecules (H_2O) are not shown for clarity.




(a)




(b)



(c)



(d)



(e)

- Q8. Given three separate solutions containing equal concentrations of formic acid ($K_a = 1.7 \times 10^{-4}$), phenol ($K_a = 1.3 \times 10^{-10}$), and acetic acid ($K_a = 1.8 \times 10^{-5}$), select the response below that has the acids arranged in order of **increasing** percent dissociation at equilibrium.
- A) formic < phenol < acetic
 B) formic < acetic < phenol
 C) acetic < formic < phenol
 D) phenol < acetic < formic
- Q9. If the concentration of hydroxide ion is 1.8×10^{-4} M in an aqueous solution at 25°C , what is the pH of the solution?
- A) 2.74
 B) 3.74
 C) 9.26
 D) 10.26
- Q10. Which solution has the largest pOH at 25°C : 0.100 M $\text{NaOH}(\text{aq})$, 0.100 M $\text{Sr}(\text{OH})_2(\text{aq})$, or 0.100 M $\text{HCl}(\text{aq})$?
- A) 0.100 M $\text{NaOH}(\text{aq})$
 B) 0.100 M $\text{Sr}(\text{OH})_2(\text{aq})$
 C) 0.100 M $\text{HCl}(\text{aq})$
 D) Impossible to determine
- Q11. What is the pH of 0.25 M $\text{LiOH}(\text{aq})$?
- A) 0.60
 B) 1.20
 C) 10.25
 D) 13.40
- Q12. A Lewis base is a(n):
- A) electron-pair donor
 B) proton acceptor
 C) electron-pair acceptor
 D) proton donor

- Q13. Which pair of substances will constitute a buffer when found in solution together:
- NaNO₂ / HNO₃
 - KCl / HF
 - HF / NaF
 - NaNO₂ / KNO₂
- Q14. A solution containing 0.30 M HClO(aq) ($K_a = 3.5 \times 10^{-8}$) and 0.25 M of NaClO(aq) would have a pH of:
- 7.46
 - 7.38
 - 7.54
 - 7.31
- Q15. A weak monoprotic acid (HA) has a pK_a of 3.94. If we need to prepare a buffer with a pH of 3.74, then we can say for sure that:
- [HA] > [A⁻]
 - [HA] = [A⁻] - 0.20 M
 - [HA] < [A⁻]
 - [HA] = [A⁻] + 0.20 M
- Q16. Which of the following acids would it be best to use to prepare a buffer with a pH of 4.25?
- HClO₂, $K_a = 1.1 \times 10^{-2}$
 - HNO₂, $K_a = 4.0 \times 10^{-4}$
 - HCH₂O, $K_a = 1.8 \times 10^{-4}$
 - H₂CO₃, $K_a = 4.3 \times 10^{-7}$
- Q17. Which chemical equation best corresponds to the K_{sp} reaction for calcium carbonate?
- $\text{CaCO}_3(\text{s}) \rightleftharpoons \text{Ca}^+(\text{aq}) + \text{C}^{3+}(\text{aq}) + 2\text{O}^{2-}(\text{g})$
 - $\text{CaCO}_3(\text{s}) \rightleftharpoons \text{Ca}(\text{aq}) + \text{CO}_3(\text{aq})$
 - $\text{CaCO}_3(\text{s}) \rightleftharpoons \text{Ca}^+(\text{aq}) + \text{CO}_3^-(\text{aq})$
 - $\text{CaCO}_3(\text{s}) \rightleftharpoons \text{Ca}^{2+}(\text{aq}) + \text{CO}_3^{2-}(\text{aq})$
- Q18. Which substance will have the **smallest** molar solubility: BaSO₄ ($K_{sp} = 1.07 \times 10^{-10}$), CaSO₄ ($K_{sp} = 7.10 \times 10^{-5}$), or FeS ($K_{sp} = 3.72 \times 10^{-19}$).
- BaSO₄
 - CaSO₄
 - FeS
 - Impossible to determine
- Q19. In which solution would ammonium fluoride be the **most soluble**?
- 0.25 M NH₄Br(aq)
 - 0.10 M NH₄Cl(aq)
 - 0.50 M LiF(aq)
 - 0.35 M KBr(aq)
- Q20. The solubility product constant for magnesium hydroxide, Mg(OH)₂ is 2.06×10^{-13} . Which solution would form a precipitate immediately upon mixing?
- A solution containing 1.0×10^{-6} M Mg²⁺ and 1.0×10^{-3} M OH⁻
 - A solution containing 1.0×10^{-5} M Mg²⁺ and 1.0×10^{-4} M OH⁻
 - A solution containing 1.0×10^{-4} M Mg²⁺ and 1.0×10^{-5} M OH⁻
 - A solution containing 1.0×10^{-3} M Mg²⁺ and 1.0×10^{-6} M OH⁻

Short Response.

Show ALL work to receive credit.

Q21. [15 pts.] (a) The pH of 0.13 M HF(aq) is 2.01. Use this information to determine K_a for HF. Be sure to write an ICE chart as part of your answer.

(b) Using your calculated K_a , predict the pH of 1.3 M HF(aq).

(c) What is the percent dissociation of HF in part (a) vs. part (b)?

Part (a) _____ %

Part (b) _____ %

Q22. [15 pts.] **(Be sure to write all relevant chemical equations and ICE/ICF charts for this question.)**
250.0 mL of 0.400 M $\text{HNO}_2(\text{aq})$, $K_a = 4.0 \times 10^{-4}$, is mixed with 50.0 mL of 0.800 M $\text{NaOH}(\text{aq})$.

(a) Calculate the pH of the solution formed.

(b) If 0.010 mol HCl is added to the solution formed in part (a), what will the final pH be?



Q23. [15 pts.] Calculate the molar solubility of aluminum hydroxide in pure water vs. in 0.200 M KOH(aq). Comment on the difference. K_{sp} for $\text{Al}(\text{OH})_3$ is 2.3×10^{-8} .

Be sure to write the K_{sp} chemical equation and an ICE chart as part of your answer! Explain any assumptions you are making.



Q24. [15 pts.] (a) Write out the chemical equation (reaction) corresponding to K_b for $\text{CH}_3\text{CH}_2\text{NH}_2(\text{aq})$, ethylamine.

(b) Write out the chemical equation (reaction) corresponding to K_w .

(c) Without performing a calculation, explain how the pH of 0.100 M $\text{NH}_3(\text{aq})$ compares with 0.100 M $\text{LiOH}(\text{aq})$.

(d) Calculate the pH of 12.0 M $\text{HCl}(\text{aq})$ (concentrated hydrochloric acid).

Bonus question

Predict whether the following aqueous salts will be acidic, basic, neutral, or whether there is not enough information to decide.

NaF _____

$\text{Al}(\text{NO}_3)_3$ _____

LiCl _____

Periodic Table of the Elements

IA		IIA										IIIA										IVA										VA										VIA										VIIA										VIII																																																																																															
1 H 1.008	2 He 4.003																																																																																																																																																												
3 Li 6.941	4 Be 9.012																																																																																																																																																												
11 Na 22.99	12 Mg 24.31																																																																																																																																																												
19 K 39.10	20 Ca 40.08	21 Sc 44.96	22 Ti 47.87	23 V 50.94	24 Cr 52.00	25 Mn 54.94	26 Fe 55.85	27 Co 58.93	28 Ni 58.69	29 Cu 63.55	30 Zn 65.39	31 Ga 69.72	32 Ge 72.61	33 As 74.92160	34 Se 78.96	35 Br 79.90	36 Kr 83.80																																																																																																																																												
37 Rb 85.47	38 Sr 87.62	39 Y 88.91	40 Zr 91.22	41 Nb 92.91	42 Mo 95.94	43 Tc [98]	44 Ru 101.1	45 Rh 102.9	46 Pd 106.4	47 Ag 107.9	48 Cd 112.4	49 In 114.8	50 Sn 118.7	51 Sb 121.8	52 Te 127.60	53 I 126.9	54 Xe 131.3																																																																																																																																												
55 Cs 132.9	56 Ba* 137.3	71 Lu 175.0	72 Hf 178.5	73 Ta 180.9	74 W 183.8	75 Re 186.2	76 Os 190.2	77 Ir 192.2	78 Pt 195.1	79 Au 197.0	80 Hg 200.6	81 Tl 204.4	82 Pb 207.2	83 Bi 209.0	84 Po [210]	85 At [210]	86 Rn [222]																																																																																																																																												
87 Fr [223]	88 Ra** [226]	103 Lr [262]	104 Rf [261]	105 Db [262]	106 Sg [266]	107 Bh [264]	108 Hs [265]	109 Mt [268]	110 [269]	111 [272]	112 [277]																																																																																																																																																		
																		* 57 La 138.9										58 Ce 140.1										59 Pr 140.9										60 Nd 144.2										61 Pm [145]										62 Sm 150.4										63 Eu 152.0										64 Gd 157.3										65 Tb 158.9										66 Dy 162.50										67 Ho 164.9										68 Er 167.3										69 Tm 168.9										70 Yb 173.0									
																		** 89 Ac [227]										90 Th 232.0										91 Pa 231.0										92 U 238.0										93 Np [237]										94 Pu [244]										95 Am [243]										96 Cm [247]										97 Bk [247]										98 Cf [251]										99 Es [252]										100 Fm [257]										101 Md [258]										102 No [259]									

$$N_A = 6.022 \times 10^{23} \text{ mol}^{-1}$$

$$R = 8.3145 \frac{\text{J}}{\text{mol} \cdot \text{K}} = 0.08206 \frac{\text{atm} \cdot \text{L}}{\text{mol} \cdot \text{K}}$$

$$M_1 V_1 = M_2 V_2$$

$$\text{pH} = -\log_{10}[\text{H}^+]$$

$$\text{pH} + \text{pOH} = 14.00 \text{ (25 }^\circ\text{C)}$$

$$K_w = 1.0 \times 10^{-14} \text{ (25 }^\circ\text{C)}$$

$$K_a \cdot K_b = K_w$$

$$\text{pH} = \text{p}K_a + \log \frac{[\text{Base}]}{[\text{Acid}]}$$

$$\text{Given: } ax^2 + bx + c = 0, \text{ then } x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

